

National Physician Survey, 2007. National Results by FP/GP or Other Specialist, Sex, Age, and All Physicians

Q28d. Which of the following procedures do you perform as part of your practice? Please check all that apply.

Question asked of FP/GPs only	FP/GP or Other Specialist		Sex			Age Groups						All FP/GPs
	FP/GP n=3841 N=29339	Other Spec n=0 N=0	Male n=2202 N=17632	Female n=1565 N=11144	NR n=74 N=563	<35 n=357 N=2735	35-44 n=940 N=7560	45-54 n=1265 N=9922	55-64 n=908 N=6607	65+ n=319 N=2123	NR n=52 N=393	Total n=3841 N=29339
	%	%	%	%	%	%	%	%	%	%	%	%
Incise & drain abscess	76.6%	.0%	79.4%	71.9%	85.0%	86.7%	80.4%	76.5%	71.8%	65.5%	79.9%	76.6%
Insert sutures/ repair lacerations	72.0%	.0%	75.4%	66.4%	77.6%	84.4%	77.5%	70.6%	66.7%	60.1%	70.9%	72.0%
Cast fractures	32.6%	.0%	38.2%	24.0%	27.5%	49.9%	37.8%	30.7%	26.1%	22.0%	26.9%	32.6%
Cryotherapy of skin lesions	63.5%	.0%	62.7%	64.8%	62.2%	74.3%	66.4%	65.8%	57.8%	46.7%	57.5%	63.5%
Excise dermal lesions	56.4%	.0%	60.6%	49.4%	65.0%	65.7%	57.8%	57.4%	53.8%	44.8%	45.7%	56.4%
Scrape skin for fungus determination	53.2%	.0%	53.6%	52.6%	53.3%	62.3%	55.9%	52.7%	51.7%	41.2%	42.3%	53.2%
Use Wood's lamp	22.8%	.0%	23.7%	21.7%	15.9%	26.4%	21.7%	25.0%	22.2%	13.2%	26.4%	22.8%
Release subungual hematoma	69.9%	.0%	74.9%	61.7%	78.4%	68.8%	69.9%	71.3%	70.5%	62.8%	73.5%	69.9%
Drain acute paronychia	61.6%	.0%	67.6%	51.9%	63.8%	62.1%	64.0%	62.6%	59.9%	55.9%	44.8%	61.6%
Pare skin callus	59.1%	.0%	60.2%	57.2%	60.7%	60.8%	60.6%	61.1%	56.9%	50.1%	54.0%	59.1%
Infiltrate local anesthetic	75.0%	.0%	79.1%	68.2%	80.6%	83.7%	79.3%	74.5%	71.4%	63.2%	69.5%	75.0%
Remove corneal or conjunctival foreign body	61.6%	.0%	68.7%	50.4%	60.5%	63.9%	60.9%	61.5%	63.0%	55.4%	68.1%	61.6%
Remove cerumen/ syringe ear canals	83.6%	.0%	83.4%	83.9%	81.6%	86.7%	84.7%	83.9%	83.1%	75.2%	83.8%	83.6%
Cauterize nose for anterior epistaxis	55.8%	.0%	61.1%	47.4%	54.7%	60.7%	58.9%	55.9%	52.9%	47.9%	52.1%	55.8%
Remove foreign body	69.6%	.0%	74.5%	61.6%	72.7%	75.1%	70.6%	69.4%	69.1%	60.6%	73.7%	69.6%
Insert nasogastric tube	32.2%	.0%	37.9%	23.5%	24.9%	48.6%	39.9%	28.7%	26.1%	20.2%	24.8%	32.2%
Test for fecal occult blood	59.2%	.0%	60.9%	55.8%	72.5%	70.8%	59.4%	57.1%	59.0%	54.2%	57.8%	59.2%
Place transurethral catheter	32.3%	.0%	38.3%	22.5%	38.4%	38.1%	36.2%	29.6%	30.4%	28.3%	36.6%	32.3%
Cryotherapy or chemical therapy for genital warts	58.4%	.0%	56.5%	61.2%	60.7%	65.0%	60.6%	60.9%	56.1%	38.2%	57.9%	58.4%
Pap smear	75.5%	.0%	71.7%	81.8%	72.3%	80.9%	72.1%	77.6%	76.8%	67.7%	73.9%	75.5%
Low forcep	5.5%	.0%	6.1%	4.7%	3.7%	4.3%	2.5%	5.8%	9.4%	5.3%	3.5%	5.5%
Mid-forcep and rotation	.8%	.0%	1.2%	.1%	1.1%	.3%	.5%	.6%	1.2%	1.7%	.0%	.8%
Vacuum extraction	10.8%	.0%	9.3%	12.9%	15.2%	19.2%	10.8%	10.8%	9.5%	3.7%	9.4%	10.8%
Splint injured extremities	48.8%	.0%	53.7%	41.1%	47.1%	57.4%	52.8%	48.3%	45.0%	37.3%	49.4%	48.8%
Bag and mask ventilation	37.5%	.0%	41.4%	31.1%	41.0%	56.6%	42.6%	35.3%	31.6%	23.0%	36.4%	37.5%
Venipuncture	41.3%	.0%	47.4%	31.7%	39.0%	41.1%	43.6%	40.3%	40.5%	38.9%	49.5%	41.3%

(continued).

National Physician Survey, 2007. National Results by FP/GP or Other Specialist, Sex, Age, and All Physicians

Q28d. Which of the following procedures do you perform as part of your practice? Please check all that apply. (continued).

Question asked of FP/GPs only	FP/GP or Other Specialist		Sex			Age Groups						All FP/GPs
	FP/GP n=3841 N=29339	Other Spec n=0 N=0	Male n=2202 N=17632	Female n=1565 N=11144	NR n=74 N=563	<35 n=357 N=2735	35-44 n=940 N=7560	45-54 n=1265 N=9922	55-64 n=908 N=6607	65+ n=319 N=2123	NR n=52 N=393	Total n=3841 N=29339
	%	%	%	%	%	%	%	%	%	%	%	%
Subcutaneous injection	82.0%	.0%	83.7%	79.4%	78.1%	80.0%	82.8%	83.5%	81.2%	75.4%	86.9%	82.0%
Intramuscular injection	83.8%	.0%	85.2%	81.7%	84.4%	80.7%	84.4%	85.7%	83.6%	77.7%	83.8%	83.8%
Insert peripheral intravenous line in both adult and child	27.2%	.0%	32.5%	18.9%	24.4%	33.4%	31.4%	26.9%	23.1%	18.9%	23.4%	27.2%
Insert central line in adult	16.2%	.0%	19.9%	10.4%	12.7%	32.4%	24.5%	13.5%	7.6%	5.0%	15.2%	16.2%
Prep for land or air transport	18.0%	.0%	20.1%	14.9%	13.2%	34.9%	25.1%	15.5%	10.4%	7.4%	12.8%	18.0%
Acupuncture	2.8%	.0%	3.4%	1.8%	2.7%	.6%	2.7%	3.5%	2.8%	2.0%	5.7%	2.8%
Hypnosis	1.8%	.0%	2.0%	1.5%	2.2%	.1%	.7%	2.0%	2.5%	4.8%	3.3%	1.8%
None of the above	3.5%	.0%	3.3%	3.8%	2.5%	.2%	2.2%	3.5%	5.0%	8.0%	2.5%	3.5%
NR	3.2%	.0%	3.3%	3.1%	3.1%	3.6%	3.4%	3.1%	2.6%	5.0%	2.9%	3.2%

Notes:

Includes only respondents who provide patient care.

Column totals may exceed 100% as this question allowed for multiple responses.

NR=No Response.

This question was asked only of family physician/ general practitioner respondents who completed the long questionnaire.

The responding sample (size: n) has been weighted to represent the population (size: N). Further detail available at www.nationalphysiciansurvey.ca.