

National Physician Survey, 2007. Results for Clinical Specialists - Internal Medicine.

Q20. Please indicate with whom you do not collaborate. Check all that apply.

| Question asked of other specialists only | | Internal Medicine Specialists | | | | | | | | | | | |
|---|-------|-------------------------------|--------------------------|-------------------------|-------------------------|-----------------------|-----------------------|---------------------------|-------------------------|-------------------------|-------------------------|-------------------------|--------------------------|
| | | Cardio n=80 N=738 | EndoMet n=32 N=240 | Gastro n=41 N=388 | Geriat n=30 N=242 | Hema n=29 N=210 | Immun n=14 N=95 | IntMed n=213 N=1607 | MedOnc n=33 N=271 | Nephro n=53 N=435 | Respir n=67 N=584 | Rheuma n=35 N=287 | Total n=627 N=5096 |
| | | % | % | % | % | % | % | % | % | % | % | % | % |
| No collaboration with Other specialists (answered by other Specialists) | Yes | 2.9% | 2.8% | 3.5% | 3.4% | ** | ** | 2.0% | .0% | 2.4% | .8% | 12.8% | 2.6% |
| | No | 97.1% | 97.2% | 93.9% | 96.6% | ** | ** | 97.0% | 100.0% | 95.4% | 99.2% | 87.2% | 96.7% |
| | NR | .0% | .0% | 2.6% | .0% | ** | ** | 1.0% | .0% | 2.2% | .0% | .0% | .7% |
| | Total | 100.0% | 100.0% | 100.0% | 100.0% | ** | ** | 100.0% | 100.0% | 100.0% | 100.0% | 100.0% | 100.0% |

Notes:

Includes only respondents who provide patient care.

NR=No Response.

This question was asked only of other specialist respondents who completed the long questionnaire.

The responding sample (size: n) has been weighted to represent the population (size: N). Further detail available at www.nationalphysiciansurvey.ca.

** Results are suppressed when there are fewer than 30 respondents.

National Physician Survey, 2007. Results for Clinical Specialists - Internal Medicine.

Q20. Please indicate with whom you do not collaborate. Check all that apply.

| Question asked of FP/GPs and other specialists | | Internal Medicine Specialists | | | | | | | | | | | Total n=627 N=5096 |
|---|--|-------------------------------|------------------------------|-------------------------|-------------------------|-----------------------|-----------------------|---------------------------|-------------------------|-------------------------|-------------------------|-------------------------|--------------------------|
| | | Cardio n=80 N=738 | Endo Met n=32 N=240 | Gastro n=41 N=388 | Geriat n=30 N=242 | Hema n=29 N=210 | Immun n=14 N=95 | IntMed n=213 N=1607 | MedOnc n=33 N=271 | Nephro n=53 N=435 | Respir n=67 N=584 | Rheuma n=35 N=287 | |
| | | % | % | % | % | % | % | % | % | % | % | % | |
| ALL - I do not collaborate with the following | No collaboration with Family physicians | 4.6% | 20.3% | 3.5% | 4.3% | ** | ** | 5.9% | 6.6% | 3.2% | 8.0% | 12.9% | 7.0% |
| | No collaboration with Nurse practitioners | 30.8% | 34.1% | 51.6% | 21.2% | ** | ** | 31.2% | 39.9% | 23.8% | 32.5% | 63.6% | 34.5% |
| | No collaboration with Psychiatric nurses | 64.8% | 54.1% | 54.7% | 34.5% | ** | ** | 47.2% | 73.6% | 53.8% | 58.0% | 68.5% | 56.2% |
| | No collaboration with Other nurses (RN, LPN, RPN) | 12.6% | 9.3% | 9.6% | .0% | ** | ** | 8.5% | 3.6% | .0% | 10.8% | 29.1% | 9.4% |
| | No collaboration with Physician assistants | 52.4% | 55.4% | 59.9% | 50.3% | ** | ** | 48.9% | 52.7% | 46.3% | 56.9% | 72.1% | 54.0% |
| | No collaboration with Dietitians/ nutritionists | 6.5% | 6.4% | 4.8% | 4.1% | ** | ** | 8.3% | 5.0% | 1.3% | 4.7% | 29.6% | 7.6% |
| | No collaboration with Occupational therapists | 24.7% | 42.7% | 28.9% | .0% | ** | ** | 15.4% | 29.2% | 8.2% | 23.3% | 16.2% | 20.4% |
| | No collaboration with Physiotherapists | 20.0% | 34.1% | 25.8% | .0% | ** | ** | 11.2% | 26.3% | 3.9% | 7.0% | 16.9% | 15.1% |
| | No collaboration with Chiropractors | 67.2% | 62.4% | 67.0% | 51.5% | ** | ** | 55.8% | 75.2% | 64.3% | 63.8% | 69.0% | 62.8% |
| | No collaboration with Psychologists | 45.4% | 28.8% | 47.0% | 14.1% | ** | ** | 42.2% | 34.5% | 48.2% | 42.8% | 51.5% | 42.1% |
| | No collaboration with Mental health counselors | 55.6% | 43.6% | 54.1% | 45.3% | ** | ** | 46.4% | 52.4% | 58.4% | 62.4% | 72.1% | 54.1% |
| | No collaboration with Addiction counselors | 60.2% | 51.9% | 56.3% | 53.9% | ** | ** | 49.1% | 68.9% | 60.6% | 60.8% | 72.1% | 57.8% |
| | No collaboration with Social workers | 17.8% | 19.5% | 26.5% | .0% | ** | ** | 14.5% | 4.3% | .0% | 12.3% | 43.1% | 15.3% |
| | No collaboration with Pharmacists | 10.9% | 15.5% | 15.9% | .0% | ** | ** | 4.4% | .0% | 1.8% | 9.2% | 30.8% | 8.4% |
| | No collaboration with Midwives | 70.0% | 61.6% | 63.5% | 60.6% | ** | ** | 54.5% | 71.6% | 62.5% | 62.1% | 75.5% | 62.9% |
| | No collaboration with Speech-language pathologists | 52.1% | 38.7% | 34.2% | 3.4% | ** | ** | 25.5% | 63.0% | 43.8% | 36.0% | 68.6% | 38.5% |
| | No collaboration with Chiropodists | 67.1% | 46.5% | 63.5% | 30.3% | ** | ** | 48.5% | 75.2% | 56.3% | 62.1% | 54.9% | 56.8% |
| No collaboration with complementary/ alternative medicine providers | 68.7% | 65.2% | 62.9% | 59.0% | ** | ** | 58.0% | 71.5% | 66.7% | 63.8% | 68.7% | 63.8% | |
| NR to questions above | .0% | .0% | 2.6% | .0% | ** | ** | 1.0% | .0% | 2.2% | .0% | .0% | .7% | |
| No collaboration at all among all respondents | 30.0% | 28.5% | 25.2% | 31.0% | ** | ** | 35.2% | 20.0% | 28.7% | 33.0% | 19.8% | 29.9% | |

Notes:

Includes only respondents who provide patient care.

Column totals may exceed 100% as this question allowed for multiple responses.

NR=No Response.

This question was asked only of those respondents who completed the long questionnaire.

The responding sample (size: n) has been weighted to represent the population (size: N). Further detail available at www.nationalphysiciansurvey.ca.

** Results are suppressed when there are fewer than 30 respondents.